

## 2020 HKIMO Prelim Problem 20

Consider the Fibonacci sequence  $1, 1, 2, 3, 5, 8, 13, \dots$ . What are the last three digits (from left to right) of the  $2020^{\text{th}}$  term?

### Solution

**Key Word** Implicit Form for Fibonacci Sequence, Binomial Expansion, Induction, Euler's Theorem, Property of Modular Arithmetic

The implicit form for Fibonacci Sequence is known to be  $F_n = \frac{1}{\sqrt{5}} \left( \frac{1+\sqrt{5}}{2} \right)^n - \frac{1}{\sqrt{5}} \left( \frac{1-\sqrt{5}}{2} \right)^n$ . In another words, the last three digits of

$$F_{2020} = \frac{1}{\sqrt{5}} \left( \frac{1+\sqrt{5}}{2} \right)^{2020} - \frac{1}{\sqrt{5}} \left( \frac{1-\sqrt{5}}{2} \right)^{2020}$$

must be computed.

Binomial expansion could be utilized to effectively expand  $F_{2020}$ .

$$\begin{aligned} \frac{1}{\sqrt{5}} \cdot \frac{1}{2^{2020}} (1 + \sqrt{5})^{2020} &= \frac{1}{\sqrt{5}} \cdot \frac{1}{2^{2020}} \left( {}_{2020}C_0 \cdot 1^{2020} \cdot (\sqrt{5})^0 + {}_{2020}C_1 \cdot 1^{2019} \cdot (\sqrt{5})^1 + \dots + {}_{2020}C_{2020} \cdot 1^0 \cdot (\sqrt{5})^{2020} \right) \\ \frac{1}{\sqrt{5}} \cdot \frac{1}{2^{2020}} (1 - \sqrt{5})^{2020} &= \frac{1}{\sqrt{5}} \cdot \frac{1}{2^{2020}} \left( {}_{2020}C_0 \cdot 1^{2020} \cdot (\sqrt{5})^0 - {}_{2020}C_1 \cdot 1^{2019} \cdot (\sqrt{5})^1 + \dots + {}_{2020}C_{2020} \cdot 1^0 \cdot (\sqrt{5})^{2020} \right) \\ F_{2020} &= \frac{1}{\sqrt{5}} \cdot \frac{1}{2^{2020}} \cdot 2 \left( {}_{2020}C_1 \cdot 1^{2019} \cdot (\sqrt{5})^1 + {}_{2020}C_3 \cdot 1^{2017} \cdot (\sqrt{5})^3 + \dots + {}_{2020}C_{2019} \cdot 1^1 \cdot (\sqrt{5})^{2019} \right) \\ &= \frac{1}{2^{2019}} \left( {}_{2020}C_1 \cdot (\sqrt{5})^0 + {}_{2020}C_3 \cdot (\sqrt{5})^2 + \dots + {}_{2020}C_{2019} \cdot (\sqrt{5})^{2018} \right) \end{aligned}$$

$a$  represents the last three digits of the  $2020^{\text{th}}$  term in the following equation.

$$F_{2020} \equiv a \pmod{1000}$$

Because  $1000 = 2^3 \cdot 5^3$ , the condition above may be split.

$$\begin{aligned} F_{2020} &\equiv b \pmod{8} \\ F_{2020} &\equiv c \pmod{125} \end{aligned}$$

### Part I: $F_{2020} \equiv b \pmod{8}$

$$\frac{{}_{2020}C_1 \cdot (\sqrt{5})^0 + {}_{2020}C_3 \cdot (\sqrt{5})^2 + \dots + {}_{2020}C_{2019} \cdot (\sqrt{5})^{2018}}{2^{2019}} \equiv b \pmod{8}$$

**Lemma.** *A repetition of remainders occurs when the Fibonacci sequence is divided by 8.*

*Proof.* First and foremost, replace 8 with  $n$  to manifest the situation.

$$1, 1, 2, 3, 5, n, n + 5, 2n + 5, 4n + 2, 6n + 7, 11n + 1, 18n, \mathbf{19n+1}, \mathbf{37n+1}, \mathbf{56n+2}, \mathbf{74n+3}, \dots$$

By induction, because the method to form the first 12 terms repeats, the remainders also retain repetition.  $\square$

$b$  is equal to 3 because  $2020 \equiv 4 \pmod{12}$ .

**Part II:**  $F_{2020} \equiv c \pmod{125}$

$$\frac{{}_{2020}C_1 \cdot (\sqrt{5})^0 + {}_{2020}C_3 \cdot (\sqrt{5})^2 + \cdots + {}_{2020}C_{2019} \cdot (\sqrt{5})^{2018}}{2^{2019}} \equiv c \pmod{125}$$

An impulse to remove the terms with  $(\sqrt{5})^6$  or with a higher degree is created. However,  $2^{2019}$  hinders the simplification. Therefore, the terms may be rewritten.

$${}_{2020}C_1 \cdot (\sqrt{5})^0 + {}_{2020}C_3 \cdot (\sqrt{5})^2 + \cdots + {}_{2020}C_{2019} \cdot (\sqrt{5})^{2018} \equiv 2^{2019}c \pmod{125}$$

Euler's theorem may be utilized for further simplification because  $GCF(2, 125) = 1$ .

$$\begin{aligned} 2^{\varphi(125)} &\equiv 1 \pmod{125} \\ 2^{125(1-\frac{1}{5})} &\equiv 1 \pmod{125} \\ 2^{100} &\equiv 1 \pmod{125} \\ 2^{2000} &\equiv 1 \pmod{125} \\ 2^{2019} &\equiv 2^{19} \pmod{125} \\ 2^{2019} &\equiv 24 \cdot 12 \pmod{125} \\ 2^{2019} &\equiv 288 \pmod{125} \\ 2^{2019} &\equiv 38 \pmod{125} \end{aligned}$$

Therefore, the following relationships are true.

$$\begin{aligned} {}_{2020}C_1 \cdot (\sqrt{5})^0 + {}_{2020}C_3 \cdot (\sqrt{5})^2 + \cdots + {}_{2020}C_{2019} \cdot (\sqrt{5})^{2018} &\equiv 38c \pmod{125} \\ {}_{2020}C_1 + {}_{2020}C_3 \cdot 5 + {}_{2020}C_5 \cdot 5^2 &\equiv 38c \pmod{125} \\ 2020 + \frac{2020 \cdot 2019 \cdot 2018}{3 \cdot 2 \cdot 1} \cdot 5 + \frac{2020 \cdot 2019 \cdot \cdots \cdot 2015}{5!} \cdot 5^2 &\equiv 38c \pmod{125} \\ 20 + 75 + 100 &\equiv 38c \pmod{125} \\ 195 &\equiv 38c \pmod{125} \end{aligned}$$

Therefore,  $38c \equiv 195 \pmod{125}$ . Moreover, because  $GCF(38, 125) = 1$ ,  $c \equiv \frac{195+125k}{38} \pmod{125}$ . Therefore,  $F_{2020} \equiv c \equiv 15 \pmod{125}$

**Part III: Computing  $F_{2020}$**

$F_{2020} \equiv 3 \pmod{8}$  and  $F_{2020} \equiv 15 \pmod{125}$  are true.

Let  $F_{2020} = 8k + 3$ .

$$\begin{aligned} 8k + 3 &\equiv 15 \pmod{125} \\ 8k &\equiv 12 \pmod{125} \\ 8k &\equiv 512 \pmod{125} \\ k &\equiv 64 \pmod{125} \quad (\because GCF(8, 125) = 1) \end{aligned}$$

Let  $k = 125k' + 64$ .

$F_{2020} = 1000k' + 67$ , or the last three digits of the 2020<sup>th</sup> term is 515. □